

U.S. Fish and Wildlife Service
Office of Subsistence Management
Fisheries Resource Monitoring Program

Abundance and Run Timing of Adult Salmon in the Tozitna River, Alaska, 2003

Annual Report for Study 03-203

This report has been prepared to assess project progress. Review comments may not be addressed in this report, but will be incorporated into the final report for this project.

Tim Sundlov
Carl Kretsinger
Bob Karlen

Bureau of Land Management
Northern Field Office
1150 University Avenue
Fairbanks, Alaska 99709

February 2004

REPORT SUMMARY PAGES

Title: Abundance and Run Timing of Adult Salmon in the Tozitna River, Alaska, 2003

Study Number: 03-203

Investigator(s)/Affiliation(s): Bob Karlen, Carl Kretsinger, and Tim Sundlov, U.S. Department of Interior, Bureau of Land Management, Northern Field Office; Julie Roberts, Tanana Tribal Council

Geographic Area: Yukon

Information Type: Stock Status and Trends

Issue(s) Addressed: Lack of escapement and run timing data in middle Yukon River Basin tributaries for Chinook *Oncorhynchus tshawytscha* and summer chum salmon *O. keta* to support Federal subsistence fishery management

Study Cost: \$27,000

Study Duration: 1 March 2003 to 29 February 2004

Abstract: The Tozitna River project is a multi-agency study to determine escapement, run timing, and age-sex-length (ASL) composition of adult Chinook and summer chum salmon in a middle Yukon Basin tributary. A resistance board weir was operated from 23 June to 12 August 2003. High stream discharge from the periods of 2 to 6 July and 26 July to 12 August prevented counting and biological sampling; no interpolation was made for these periods. Comparison of run timing to 2002 and to other tributaries with similar run timing suggests the majority of the Chinook salmon run was counted. The escapement for Chinook salmon was 1,819 (18% female). Age groups 1.2 and 1.3 accounted for 28 and 52% of the escapement, respectively. The escapement for summer chum salmon was 8,487 (34% female). The majority of the summer chum salmon escapement was not counted and ASL composition were determined with partial escapement data. Age 0.3 comprised 86% of the chum escapement.

Key Words: Chinook salmon, chum salmon, *Oncorhynchus tshawytscha*, *O. keta*, resistance board weir, sex ratio, spawning adults, stock/status trends, subsistence fishery, Tozitna River, Yukon River drainage

Project Data: *Description* - Data for this study consist of biological samples (scales, scale impressions, fin clips, and processed DNA samples), escapement, and sampling data (date, age, sex, and length) on Chinook and summer chum salmon. *Format* - Scale impressions were created on cellulose acetate cards. Processed DNA samples are stored at -20 °C. Scale and genetic data are stored in Microsoft Access and Excel databases. *Custodians* - Genetic samples and data: U.S. Fish and Wildlife Service Fish Genetics

Laboratory (USFWS-FGL), 1011 East Tudor Road, Anchorage, Alaska 99503. Sampling data: Bureau of Land Management (BLM), Northern Field Office, 1150 University Avenue, Fairbanks, Alaska 99709 and Alaska Department of Fish and Game, Division of Commercial Fisheries (ADF&G-DCF), 333 Raspberry Road, Anchorage, Alaska 99518. Availability - Access to biological samples and data is available upon request to the custodians.

Report Availability: Please contact either the author(s) or Alaska Resources Library and Information Services to obtain a copy of this report.

Citation: Sundlov, T.J., C. F. Kretsinger and B. Karlen. 2003. Abundance and Run Timing of Adult Salmon in the Tozitna River, Alaska, 2003. USFWS Office of Subsistence Management, Fisheries Resource Monitoring Program, Annual Report No. 03-203, Anchorage, Alaska.

TABLE OF CONTENTS

Abstract.....	i
Key Words	i
Citation.....	ii
Introduction.....	1
Study Area	2
Methods.....	3
Weir and Trap	3
Biological Data	3
Abundance Downstream of the Weir.....	3
Genetic Samples.....	4
Abiotic Measurements	4
Data Analysis	4
Results.....	4
Weir and Trap Operation	4
Escapement	5
Chinook salmon	5
Summer chum salmon.....	5
Age-Sex-Length.....	5
Chinook Salmon.....	5
Summer Chum Salmon.....	6
Abiotic Measurements	6
Discussion.....	6
Acknowledgements.....	8
Literature Cited.....	8
Footnotes.....	11

Tables

Table 1. Daily and cumulative counts for Chinook and summer chum salmon with quartiles shown (25, 50, 75%) of the cumulative escapement, Tozitna River, Alaska, 2003.....	12
Table 2. Chinook salmon escapement age composition by stratum and sex, Tozitna River, Alaska, 2003.....	13
Table 3. Proportion and estimated number of female Chinook salmon, Tozitna River, Alaska, 2003.	14
Table 4. Chinook salmon mid eye to fork length (mm) by age and sex, 2003, Tozitna River, Alaska.....	14
Table 5. Summer chum salmon escapement age composition by stratum and sex, Tozitna River, Alaska, 2003.	15
Table 6. Proportion and estimated number of female summer chum salmon, Tozitna River, Alaska, 2003.....	16
Table 7. Summer chum salmon mid eye to fork length (mm) by age and sex, 2003, Tozitna River, Alaska.	16

Table 8. Comparison of preliminary Chinook salmon escapement age composition by sex at the EF of Andreafsky River, Gisasa River, Henshaw Creek, and the Tozitna River, Alaska, 2003.....	17
---	----

Figures


Figure 1. Location of the Tozitna River weir, Alaska, 2003.	18
Figure 2. Chinook salmon daily counts with quartiles shown (25, 50, and 75%) of cumulative escapement for the period 26 June – 26 July, 2003, Tozitna River, Alaska.	19
Figure 3. Daily discharge (m ³ /s) for the period 21 June – 11 August 2003, Tozitna River, Alaska.....	20
Figure 4. Map showing the locations of four weirs in Alaska monitoring Chinook salmon escapement on Yukon River tributaries: East Fork of the Andreafsky River, Henshaw Creek, Gisasa River, and the Tozitna River.....	21

INTRODUCTION

Conservation of salmon in the Yukon River drainage is complex and challenging for fisheries managers because of several biological and social factors: mixed-stocks, large geographic spawning distribution, overlapping and compressed run timing, recent declines in escapement, multiple user groups, and multi-agency management. Several plans and policies have been created to manage the Yukon River salmon escapement (see Holder and Senecal-Albrecht 1998). Mostly, the Yukon River salmon escapement is managed based on sustained yield, defined as the average annual yield resulting from an escapement level that can be maintained on a continuing basis.

In 1998, the Yukon River Comprehensive Salmon Plan for Alaska (YRCSPA) was developed to improve salmon management in the Yukon Area. On October 1, 1999, the Federal government joined the State of Alaska in managing Yukon River fisheries, assuming responsibility for subsistence fisheries management in inland navigable waters on, and adjacent to, Federal conservation lands (Buklis 2002).

In 2000, BLM in Alaska received a Congressional appropriation for Yukon River salmon restoration. In response to this appropriation, the BLM convened interagency coordination meetings to determine the most beneficial use of the funding. Emphasis was placed on funding projects that would satisfy both the BLM and Yukon River fisheries management. Yukon River fisheries managers placed a priority on addressing escapement and run timing data gaps in the middle Yukon River Sub-Basin for Chinook *Oncorhynchus tshawytscha* and summer chum *O. keta* salmon, as identified in the YRCSPA (Holder and Senecal-Albrecht 1998). After interagency coordination meetings, the BLM chose the Tozitna River. The BLM had in 1986 designated the Tozitna River an Area of Critical Environmental Concern for the protection of salmon spawning habitat and had identified acquisition of baseline resource data as a management objective (BLM 1986; Knapman 1989). In addition to addressing data gaps identified in the YRCSPA, salmon escapement and run timing data collected on the Tozitna River would assist the BLM in fulfilling its management objectives.

Accurate escapement estimates from spawning tributaries are an important fisheries management tool used to assist in the determination of duction, marine survival, harvest, and spawner recruit relationships (Neilson and Green 1981; Labelle 1994). Although aerial surveys escapement estimates on the Tozitna River have been conducted by ADF&G since 1959, results of aerial surveys are inherently variable (Schultz et al. 1993) and should only be used to examine trends in relative escapement abundance (Barton 1984). Samples taken at weirs are considered to be the least biased and most accurate data available for assessing escapement and age composition of a mixed stock fishery (Halupka et al. 2000).

To accurately assess escapement of Chinook and summer chum in the Middle Yukon Sub-Basin, the BLM has operated a resistance board weir on the Tozitna River since 2002. Objectives of the project are to:

- (1) Determine escapement of Chinook and summer chum salmon;
- (2) Describe the run timing of Chinook and summer chum salmon;
- (3) Estimate relative abundance of Chinook and summer chum salmon downstream of the weir and document spawning locations using aerial survey techniques; and
- (4) Estimate age-sex-length composition of Chinook and summer chum salmon such that simultaneous 90% confidence intervals have maximum width of 0.20.

Additional project tasks are to:

- (1) Measure water temperature, turbidity, precipitation, stream height, and determine stream discharge;
- (2) Collect Chinook salmon fin tissue samples for the USFWS genetic stock analysis; and
- (3) Recover radio telemetry tags for the Yukon Basin Chinook radio telemetry study.

STUDY AREA

The Tozitna River is a large, clear-water, northern tributary to the middle Yukon River, with a watershed area of 4,212 km², 90% of which the BLM manages (Figure 1). The watershed originates in the southeastern Ray Mountains at 1,676 m and flows southwesterly approximately 207 km to its confluence with the Yukon River (1,096 river km), 16 km west of Tanana. The average yearly precipitation is 32 cm⁽¹⁾ with 62% occurring between June and September. Average monthly ambient temperature ranges from -28 to 22 °C⁽¹⁾. The river is usually ice-free in May, and freeze-up commonly occurs by November (J. Blume, Tozitna River homesteader, Fairbanks, personal communication). Peak discharge is correlated with spring snowmelt or high-intensity rainstorms during the summer. Water turbidity remains low for the period from late June through early August, except for periods of high-intensity precipitation. Fish species in the Tozitna River include Chinook salmon, summer and fall chum salmon (Barton, 1984), coho salmon *O. kisutch*, sockeye salmon *O. nerka*, Dolly Varden *Salvelinus malma*, Arctic grayling *Thymallus arcticus*, northern pike *Esox lucius*, burbot *Lota lota*, round whitefish *Prosopium cylindraceum*, slimy sculpin *Cottus cognatus*, and longnose sucker *Catostomus catostomus*.

The weir site is approximately 80 km upstream from the mouth of the Tozitna River. The weir is located between a downstream riffle and upstream deep meander pool. At this location the average wetted width at summer flows is 64 m with an average depth of 0.7 m. This site is downstream of most Chinook salmon spawning (Kretsinger and Sundlov 2001, in preparation). The cross section is gradually sloping and the substrate consists of sand to cobble.

METHODS

Weir and Trap

Salmon escapement, run timing, and composition were assessed by counting and sampling fish as they passed through the resistance board weir fitted with an in-stream live trap. Construction and installation of the weir were as described by Tobin (1994). The trap was preconstructed (Mackey Lake Co., Soldotna, AK) and incorporated into the weir on the upstream side. The weir was 65 m in width and was operational on 19 June. The weir panels, base rail, and trap were visually inspected daily for possible escapement openings and for removal of accumulated debris.

Biological Data

All salmon passing through the weir and live trap were counted and identified to species. Observers wore polarized sunglasses to facilitate in fish identification. The counting schedule was 24 h/d, 7 d/wk and consisted of one observer for each 6 h period. Except for sampling, salmon were counted without migration interruption as they proceeded through the open trap. Hourly counts were summed to achieve a daily count (0000 – 2359 hours). No interpolation was made for missed counting periods during high flows, which occurred from 2 to 6 July and 26 July to 12 August. Run timing was calculated by the proportion of daily to cumulative passage to determine quartile (25, 50, and 75%) dates and peak and median date of passage.

The live trap was used to capture salmon for biological sampling. The upstream gate of the trap was closed for periods to obtain an adequate sample size. During sampling, a dip-net was used to capture salmon in the live trap. Salmon were then placed in a submerged, aluminum cradle for identifying species and sex, measuring, and removing scale(s) and a fin clip. Lengths were measured to the nearest 5 mm from mid eye to fork of the caudal fin. Morphological maturation characteristics were used to determine sex. One scale for chum and three scales for Chinook salmon were removed from the left side, two rows above the lateral line and on a diagonal line from the posterior end of the dorsal fin to the anterior end of the anal fin (Anas 1963; Mosher 1968). Scales were then placed on numbered gum cards and sent to ADF&G-DCF in Anchorage for aging. Aging was conducted by creating impressions on cellulose acetate cards with a heated hydraulic press (Clutter and Whitsel 1956) and then examining the scale annuli patterns (Gilbert 1922). European notation (Koo 1962) was used to record the ages. A holding pen (4 m x 2 m) was constructed adjacent to the trap, and after sampling, fish were transferred and held for 0.5 h. The holding pen provided sampling recovery with low water velocity that facilitated upstream migration.

Abundance Downstream of the Weir

To estimate relative abundance of Chinook and summer chum salmon downstream of the weir and document spawning locations, an aerial survey was to be conducted late in the run, but high stream flows and turbidities prevented this from occurring.

Genetic Samples

Throughout the run, dorsal fins were clipped from 250 Chinook and 250 summer chum salmon to provide tissue samples for genetic analysis. Fin clips were placed in 2 ml sample vials filled with 95% ethanol and sent to USFWS-CGL, Anchorage for processing.

Abiotic Measurements

Water temperature, turbidity, precipitation, and stream height data were collected daily from the period 21 June to 11 August. Water temperature was monitored with an Onset® Tidbit temperature logger at the weir, recorded every hour, and summarized as daily mean. Turbidity was measured using a HACH model 2008 Turbidimeter. Precipitation was measured daily for the previous 24 h with a rain gauge. A staff gauge was surveyed to reference marks at the weir to record relative stream height.

To determine stream discharge, water velocity was measured over a range of staff gauge heights using a Price AA current meter. Stream height measurements were used as the independent variable to estimate stream discharge for days when discharge was not measured. A stream height versus discharge rating was developed by combining the direct discharge measurements and computer-simulated peak flows using log-log regression (Rantz et al. 1982).

Data Analysis

Temporally stratified random sampling design (Cochran 1977) was used to collect and analyze ASL data, with statistical weeks defining strata. Sample size goals were established so that simultaneous 90% interval estimates of sex and age composition for each week have maximum widths of 0.20 (Bromaghin 1993). Strata began on Monday and ended the following Sunday with a weekly sample size target of 154 chum and 169 Chinook salmon sampled uniformly throughout the week (25 fish/species/day). All target species within the trap at the time of sampling were sampled to avoid bias. The first and last sampling strata are greater than a week because of low escapement for those periods.

RESULTS

Weir and Trap Operation

Weather systems in the summer often bring periods of rain to the interior of Alaska and occasionally produce high stream discharge, which can submerge weir panels and allow salmon to migrate over the weir undetected. Two strong precipitation events occurred at the weir in early and late July (Figure 3). During the period of 2 to 3 July, 2.1 cm of rain was recorded. This event resulted in submerged weir panels from 2 to 6 July, allowing salmon to migrate over the weir undetected. Large trees were caught on the weir panels, base rail, and trap during this period. Entire trees were pulled over the weir and pushed off the trap and base rail when possible, but on occasion debris had to be separated into smaller pieces and removed. On 3 July, a large tree and its root mass was caught by the trap, perpendicular to the current, and generated enough force to cause the trap's earth

anchors to fail; subsequently the trap was washed 300 m downstream. The trap broke several pickets in four panels as it moved downstream over the weir. The trap was then winched upstream back into position and the weir panels were repaired. The weir was back in operation on 6 July at 1830 hours. Another strong system in late July brought record rainfall to the central interior. The high stream discharge submerged the panels from 26 July to 7 August, a result of 5.8 cm of rain. The weir and trap remained in place, but several trees had damaged the trap. On 8 August it was assumed from previous run timing data (Kretsinger and Sundlov 2001, in preparation), that the migration period of summer chum and Chinook salmon was ending, and removal of the weir and trap was initiated during an available window of reduced stream discharge. The removal was completed on 12 August.

Escapement

Chinook Salmon

Chinook salmon (N = 1,819) passed through the weir from 26 June to 26 July (Table 1). Daily Chinook escapement for the last four complete days of counting was < 3% of the cumulative escapement. Gisasa River Chinook have similar run timing (Kretsinger and Sundlov 2001, in preparation), and 2003 preliminary data indicate that Gisasa River midpoint was 13 July, a day earlier than the Tozitna River. More than 95% of the Gisasa River Chinook salmon cumulative escapement had occurred by 25 July, suggesting that the majority of the Tozitna River Chinook run was counted. The quartile days (25, 50, and 75%) of cumulative passage for Chinook salmon were 9, 14, and 19 July, respectively (Table 1 and Figure 2). Interpolation of Chinook salmon escapement for the missed counting period of 2 to 6 July did not affect run timing date determination. The date of peak passage was 9 July (n = 365), and the 7 d period between 9 and 15 July accounted for 53% of the escapement. The midpoint date of passage for Chinook salmon in the lower Yukon River was six days earlier than the average date for the midpoint (ADF&G 2003). The Tozitna River escapement midpoint was six days earlier than 2002.

Summer Chum Salmon

Summer chum salmon (N = 8,487) migrated through the weir from 25 June to 26 July (Table 1). Determinations of run timing and escapement for summer chum salmon were not possible because a significant portion of the run was missed due to high stream discharge.

Age-Sex-Length

Chinook Salmon

The sex composition of Chinook salmon was 18% female, ranging from 3 to 55% throughout weekly sampling stratum (Table 3). Overall, Chinook salmon were predominantly age 1.3 (51.7%) and 1.2 (27.7%) (Table 2). Females were generally older (68% age 1.4 and only 31% age 1.3) than males (56% age 1.3 and 34% age 1.2). The age structure of the run was reflected in size, with females ranging from 725mm to 950mm (97% \geq 750mm) and the smaller males ranging from 330mm to 975mm (with 18% \geq

750mm) (Table 4). Mean length of females age 1.3 and 1.4 was greater than that of same-age males.

Summer Chum Salmon

The sex composition of summer chum salmon was 34% female, ranging from 0 to 37% throughout weekly sampling stratum (Table 6). Overall, chum salmon were predominantly age 0.3 (86%) (Table 5). Female chum salmon ranged from 500 to 640 mm and male chum salmon ranged from 510 to 675 mm (Table 7).

Abiotic Measurements

Water temperature (°C) at the weir ranged from 6.1 to 14.0 and averaged 9.6. Turbidity (NTU) ranged from 0.8 to 56.9 and averaged 3.8. Total precipitation for the period was 10.2 cm. Stream height (cm) fluctuated from 98 to 207 and averaged 130. Daily discharge (m³/s) ranged from 4 to 83 and averaged 18 (Figure 3).

DISCUSSION

Assessment of sex and age composition of salmon stocks offers insight into the effects of fisheries management regulations and environmental influences. In 2003, there were four weirs in Alaska monitoring Chinook salmon escapement on Yukon River tributaries: East Fork of the Andreafsky River, Henshaw Creek, Gisasa River, and the Tozitna River (Figure 4). These weirs provide the most accurate method for assessing sex and age composition. This section discusses the low proportion of females in escapement data at these four weirs, following the conclusion in ADG&G (2002) and Wiswar (2001) that the proportion of females should be taken into account when assessing escapement.

The Tozitna River had the lowest proportion of female Chinook salmon of the four Yukon River tributary weirs in 2002 (Table 8) and 2003, at 13% and 18%, respectively. Although the 2003 Tozitna River Chinook escapement was 21% above that of 2002, the proportion of females was only 5% higher. The low proportion of female Chinook salmon is not unique to the Tozitna River and has been documented for other Yukon River tributaries. From 1994 - 2000, the Gisasa River averaged 30% female, ranging from 17 - 42%, and the East Fork of the Andreafsky River from 1994 - 1998 averaged 35% female, ranging from 25 - 42% (Wiswar 2001, Tobin and Harper 1999). In 2003, preliminary data indicate that three of the four Yukon tributary weirs had sex ratios favoring males (61 - 82%) (Table 8).

Recently, there has been speculation that the disease-induced mortality caused by the internal parasite *Ichthyophonous hofei* has played a role in the selective mortality of female Chinook salmon in the Yukon River. Kocan et al. (2003) reported that significantly more Yukon River females than males were infected during 1999 - 2002. However, in 2003 the infection in females was not significantly different from males,

indicating this may not be a plausible explanation for the low number of female Chinook salmon.

Low female Chinook sex ratios at weir escapement projects are largely the result of the low proportion of age 1.4 females, the predominate age class among females (Harper and Watry 2001). In 2003, preliminary data indicate age 1.4 female Chinook salmon represented 12 – 29% of escapement at the four Yukon tributary weirs (Table 8) and 39% in the lower commercial harvest. The Tozitna River escapement in 2003 had the lowest proportion of age 1.4 female Chinook at 12%. ASL composition is not available for the subsistence harvest, although in the lower Yukon River it is assumed to be similar to the lower commercial harvest, since the same gear is used (Menard 1996).

Chinook salmon harvest in the Yukon River is comprised predominately of commercial and subsistence gillnet fisheries. In 2003, the majority of commercial harvest (91%) occurred in the lower Yukon River, where gillnet mesh size (stretched) was restricted to eight inch or greater (ADF&G 2003). Fully 87% of the Chinook salmon subsistence harvest in 2002 was taken with gillnets, and 44% of the harvest occurred in the lower Yukon River (0 - 311 km) (Brase 2003). There were no gillnet mesh restrictions for the subsistence fishery in 2003, although it is thought the majority of Yukon River gillnet subsistence fishers use eight inch or greater because this is a requirement needed to participate in the commercial fishery.

Yukon Chinook populations are heterogeneous in age, size, and sex, and all individuals are not equally vulnerable to harvest. For example, Tozitna River female Chinook exhibited sexual dimorphism, with females longer than males of the same age. Large mesh gillnets used during unrestricted mesh-size openings select older, larger Chinook salmon, which include a much larger proportion of females than small mesh-size periods (ADF&G 2002). Salmon encounter gillnets from the Yukon River mouth to river km 493 with cumulative harvest effects, as the largest and oldest fish are continuously selected as they migrate upstream. The Tozitna River escapement project is the furthest upstream of the four Yukon River tributary weirs, and in 2002 and 2003 it had the lowest female sex ratio and the lowest proportion of age 1.4 female Chinook salmon (Table 8).

Another possible explanation of low abundance of age 1.4 female Chinook is their possible differential exposure to ocean mortality. The average age of maturity for Yukon River Chinook is 6.12 years for females and 5.64 years for males (McBride et al. 1983). Therefore, female's longer duration of ocean residency may increase exposure to mortality.

Preliminary results indicate that the selective harvest of larger salmon in the commercial and subsistence fisheries and/or differential mortality may have contributed to the low proportion of female Chinook salmon in the Tozitna River escapement. Reduction and removal of the largest and potentially most successful spawners reduces the overall fitness of a population and reduces the ability to compensate for environmental and anthropogenic impacts (Livingston 1998). Furthermore, Ricker (1981) argues that selective harvest by fisheries changes the genetic basis for maturation and can result in a

reduction in adult size. Although long-term weir escapement data is not available on the Tozitna River and there is no conclusive data for selective harvest and/or differential mortality, the low proportion of returning females warrants further evaluation and ongoing monitoring.

ACKNOWLEDGEMENTS

The authors are grateful to the following individuals for providing data collection and assistance under sometimes challenging field conditions: Harris Hyslop, Roxanne Kennedy, Arnold Marks, and Daniel Sam, Tanana Tribal Council, Tanana, Alaska; Seth Beaudreault, Matthew Churchman, Kelly Hoover, and Arthur Niven, Student Conservation Association; Tom Edgerton, Steve Lundeen, Craig McCaa, Jason Post (relief crew leader), Sage, and Chad Thompson, Bureau of Land Management, Fairbanks; Kimberly Elkin and Valli Peterson, Tanana Chiefs Conference, Fairbanks; Cliff Schleusner and Beth Spangler, U.S. Fish and Wildlife Service, Anchorage; Josh Holbrooke, Talkeetna, Alaska. We also appreciate the assistance of the Alaska Department of Fish and Game, Commercial Fisheries Division, Anchorage in the scale sampling analysis and Julie Roberts of the Tanana Tribal Council for her assistance with the cooperative agreement. A special thanks goes to Mr. Jack Blume for the use of his private airstrip on the Tozitna River. Thanks also to Fred Anderson, Taylor Brelsford, David Esse, Craig McCaa, Ingrid McSweeney, Robin Mills, Dave Parker, Cliff Schleusner, and Dennis Tol for their helpful reviews. The U.S. Fish and Wildlife Service, Office of Subsistence Management, Anchorage, provided \$27,000 in funding through the Fisheries Resource Monitoring Program under FWS agreement number 70181-3-N163. Additional funding was provided by the Bureau of Land Management, Northern Field Office, Fairbanks.

LITERATURE CITED

- Anas, R.E. 1963. Red salmon scale studies. Pages 114-116 in International North Pacific Fisheries Commission Annual Report, 1961. Vancouver, British Columbia.
- ADF&G (Alaska Department of Fish and Game). 2002. 2002 Yukon Area subsistence, personal use, and commercial salmon fisheries outlook and management strategies. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report Number 3A02-35, Anchorage, Alaska.

- ADF&G (Alaska Department of Fish and Game). 2003. Preliminary 2003 Yukon Area chinook and summer chum salmon fishery summary. Alaska Department of Fish and Game, Division of Commercial Fisheries, Anchorage, Alaska.
- Barton, L.H. 1984. A catalog of Yukon River salmon spawning escapement surveys. Alaska Department of Fish and Game, Technical Data Report 121, Juneau, Alaska.
- Brase, A. L. J., and H.H. Hamner. 2003. Subsistence and personal use salmon harvests in the Alaska portion of the Yukon River drainage, 2002. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report Number 3A03-13, Anchorage, Alaska.
- Bromaghin, J.F. 1993. Sample size determination for interval estimation of multinomial probabilities. *The American Statistician* 47(3):203-206.
- Buklis, L. S. 2002. Subsistence fisheries management of federal public lands in Alaska. *Fisheries* 27(7):10-18.
- BLM (Bureau of Land Management). 1986. Resource management plan and record of decision for the Central Yukon planning area. Bureau of Land Management, Kobuk District Office, Document Number BLM-AK-PT-86-031-1610-026, FF085208, Fairbanks, Alaska.
- Clutter, R.I., and S.E. Whitsel. 1956. Collection and interpretation of sockeye salmon scales. *Bulletin of the International Pacific Salmon Fisheries Commission* No. 9. Westminister, BC, Canada.
- Crane, P.A., W.J. Spearman and L.W. Seeb. 2001. Yukon River chum salmon: Report for genetic stock identification studies, 1992-1997. Alaska Department of Fish and Game, Regional Information Report Number 5J01-08, Juneau, Alaska.
- Gilbert, C.H. 1922. The salmon of the Yukon River. Bureau of Fisheries, *Bulletin* 38:317-322, Washington.
- Halupka, K.C., M.D. Bryant, M.F. Wilson, and F.H. Everest. 2000. Biological characteristics and population status of anadromous salmon in Southeast Alaska. General Technical Report: PNW-GTR-468, U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station, Juneau, Alaska.
- Harper, K.C., and C. B. Watry. 2001. Abundance and run timing of adult salmon in the Kwethluk River, Yukon Delta National Wildlife Refuge, Alaska, 2000. U.S. Fish and Wildlife Service, Kenai Fishery Resources Office, Fishery Data Series Number 2001-4, Kenai, Alaska.

- Holder, R.R., and D. Senecal-Albrecht (compilers). 1998. Yukon River comprehensive salmon plan for Alaska. Alaska Department of Fish and Game, Juneau, Alaska.
- Knapman, L.N. 1989. Watershed activity plan for the Tozitna River watershed Area of Critical Environmental Concern. Bureau of Land Management, Kobuk District Office, Document Number BLM-AK-PT-89-013-7200-070, Fairbanks, Alaska.
- Kocan, R., P. Hershberger, and J. Winton. 2003. Effects of *Ichthyophonus* on survival and reproductive success of Yukon River Chinook salmon. Federal Subsistence Fishery Monitoring Program, Final Project Report Number FIS 01-200. U.S. Fish and Wildlife Service, Office of Subsistence Management, Fishery Information Services Division, Anchorage, Alaska.
- Koo, T.S.Y. 1962. Age designation in salmon. Pages 37-48 in T.S.Y. Koo, editor. Studies of Alaska red salmon. University of Washington Publications in Fisheries, New Series, Volume I, Seattle, Washington.
- Kretsinger, C. F., and T.J. Sundlov. *In preparation*. Abundance and run timing of adult salmon, with observations of streamflow and water quality, in the Tozitna River Area of Critical Environmental Concern, Alaska 2001. Bureau of Land Management, Northern Field Office, Fairbanks Alaska.
- Labelle, M. 1994. A likelihood method for estimating Pacific salmon based on fence counts and mark-recapture data. Canadian Journal of Fisheries and Aquatic Sciences 51:552-556.
- Livingston, P.A. (editor). 1998. Draft Bering Sea ecosystem research plan. Alaska Fisheries Science Center, Seattle, WA.
- McBride, D.N. and S.L. Marshall. 1983. Feasibility of scale pattern analysis to identify the origins of chinook salmon (*Oncorhynchus tshawytscha* Walbaum) in the Lower Yukon River commercial gillnet fishery, 1980-1981. Alaska Department of Fish and Game, Division of Commercial Fisheries, Informational Leaflet 208, Juneau, Alaska.
- McBride, D.N., H.H. Marshall, and L.S. Bulkis. 1983. Age, sex and size of Yukon River salmon catch and escapement, 1982. Alaska Department of Fish and Game, Technical Data Report 90, Juneau, Alaska.
- Menard, J. 1996. Age, sex and length of Yukon River salmon catches and escapements, 1994. Alaska Department of Fish and Game, Commercial Fisheries Management and Development Division, Regional Information Report Number 3A96-16, Anchorage, Alaska.
- Mosher, K. 1968. Photographic atlas of sockeye salmon scales. Fishery Bulletin 67:243-280.

- Neilson, J.D., and G.H. Geen. 1981. Enumeration of spawning salmon from spawner residence time and aerial counts. *Transactions of the American Fisheries Society* 110:554-556.
- Rantz, S.E., et al. 1982. Measurement and computation of streamflow, volume 1 and 2 of Water-Supply Paper 2175. U.S. Geological Survey, Washington, DC.
- Ricker, W.E. 1981. Changes in the average size and average age of Pacific salmon. *Canadian Journal of Fisheries and Aquatic Sciences*. 38:1636-1656.
- Schultz, K.C., R.R. Holder, L.H. Barton, D.J. Bergstrom, C. Blaney, G.J. Sandone, and D.J. Schneiderhan. 1993. Annual management report for subsistence, personal use, and commercial fisheries of the Yukon area, 1992. Alaska Department of Fish and Game, Regional Information Report Number 3A93-10, Anchorage, Alaska.
- Tobin, J.H. 1994. Construction and performance of a portable resistance board weir for counting migrating adult salmon in rivers. U.S. Fish and Wildlife Service, Kenai Fishery Resource Office, Alaska Technical Report Number 22, Kenai, Alaska.
- Tobin, J.H. III, and K.C. Harper. 1999. Abundance and run timing of adult salmon in the East Fork Andreafsky River, Yukon Delta National Wildlife Refuge, Alaska, 1998. US. Fish and Wildlife Service, Kenai Fishery Resources Office, Alaska Fishery Data Series Number 99-3, Kenai, Alaska.
- Wiswar, D. W. 2001. Abundance and run timing of adult salmon in the Gisasa River, Koyukuk National Wildlife Refuge, Alaska, 2000. U.S. Fish and Wildlife Service, Fairbanks Fishery Resource, Alaska Fisheries Data Series Number 2001-1, Fairbanks, Alaska.

FOOTNOTES

¹ 1949 – 2003 average monthly temperature and precipitation data for the Tanana FAA Airport, Alaska, supplied by Western Regional Climate Center, Reno, Nevada.

Table 1. Daily and cumulative counts for Chinook and summer chum salmon with quartiles shown (25, 50, and 75%) of cumulative escapement, Tozitna River, Alaska, 2003.

Date	Chinook			Summer chum	
	Daily Count	Cumu Count	Cumu Proportion	Daily Count	Cumu Count
6/25	0	0	0.00	2	2
6/26	2	2	0.00	0	2
6/27	1	3	0.00	2	4
6/28	0	3	0.00	0	4
6/29	0	3	0.00	2	6
6/30	1	4	0.00	1	7
7/1	0	4	0.00	0	7
7/2 ^a	4	8	0.00	1	8
7/3 ^a	0	8	0.00	0	8
7/4 ^a	0	8	0.00	0	8
7/5 ^a	0	8	0.00	0	8
7/6 ^a	25	33	0.02	3	11
7/7 ^a	21	54	0.03	3	14
7/8	52	106	0.06	13	27
7/9	365	471	0.26	92	119
7/10	140	611	0.34	146	265
7/11	50	661	0.36	106	371
7/12	90	751	0.41	138	509
7/13	153	904	0.50	72	581
7/14	62	966	0.53	155	736
7/15	98	1064	0.58	184	920
7/16	61	1125	0.62	72	992
7/17	46	1171	0.64	65	1057
7/18	166	1337	0.74	238	1295
7/19	123	1460	0.80	472	1767
7/20	92	1552	0.85	741	2508
7/21	117	1669	0.92	864	3372
7/22	25	1694	0.93	458	3830
7/23	12	1706	0.94	829	4659
7/24	54	1760	0.97	1403	6062
7/25	46	1806	0.99	1689	7751
7/26 ^a	13	1819	1.00	736	8487

Boxed areas = quartiles (25, 50, and 75%)

Cumu = Cumulative

^aWeir was not operational due to high stream discharge.

Table 2. Chinook salmon escapement age composition by stratum and sex, Tozitna River, Alaska, 2003. Standard error in parentheses.

		Brood Year and Age												
Strata	Run	Sex	2000		1999		1998		1997		1996		Total	
			1.1		1.2		1.3		1.4		1.5		N	%
			N	%	N	%	N	%	N	%	N	%	N	%
6/26-7/7	54	Males	0	0.0	9 (7.2)	23.7	22 (8.2)	57.9	6 (6.1)	15.8	0	0.0	37	97
		Females	0	0.0	0	0.0	1 ^a	2.6	0	0.0	0	0.0	1	3
		Subtotal	0	0.0	9 (7)	23.7	23 (8)	60.5	6 (6)	15.8	0	0.0	38	100
7/8-7/14	912	Males	1 (.5)	0.5	61 (3.5)	29.3	109 (3.6)	52.4	13 (1.9)	6.3	0	0.0	184	89
		Females	0	0.0	0	0.0	12 (10.4)	5.8	12 (10.4)	5.8	0	0.0	24	12
		Subtotal	1 (.5)	0.5	61 (3.2)	29.3	121 (3.4)	58.2	25 (2.3)	12.1	0	0.0	208	100
7/15-7/21	703	Males	1 (.6)	0.5	61 (3.7)	28.6	92 (3.8)	43.2	15 (2.2)	7.0	1 (0.6)	0.5	170	80
		Females	0	0.0	0	0.0	11 (6.7)	5.2	31 (6.9)	14.6	1 (2.3)	0.5	43	20
		Subtotal	1 (.5)	0.5	61 (3.1)	28.6	103 (3.4)	48.4	46 (2.8)	21.6	2 (0.7)	1.0	213	100
7/22-7/28	150	Males	0	0.0	8 (11.6)	19.0	8 (11.6)	19.0	3 (8.6)	7.1	0	0.0	19	45
		Females	0	0.0	0	0.0	4 (8.1)	9.5	19 (8.1)	45.2	0	0.0	23	55
		Subtotal	0	0.0	8 (6.1)	19.0	12 (7.1)	28.5	22 (7.8)	52.3	0	0.0	42	100
Subtotal		Males	2 (.5)	0.4	139 (4.9)	27.7	231 (5.0)	46.1	37 (3.3)	7.4	1 (0.4)	0.2	410	82
		Females	0	0.0	0	0.0	28 (8.8)	5.6	62 (8.9)	12.4	1 (1.4)	0.2	91	18
Total	1819		2	0.4	139 (3.6)	27.7	259 (4.1)	51.7	99 (3.4)	19.8	2 (0.4)	0.4	501	100

^aStandard error was not calculated because of low sample size.

Table 3. Proportion and estimated number of female Chinook salmon, Tozitna River, Alaska, 2003.

Strata	Run	N	Percent Female	Estimated # Females
6/26-7/7	54	38	2.6 (2.2)	1
7/8-7/14	912	208	11.5 (1.1)	105
7/15-7/21	703	213	20.2 (1.5)	142
7/22-7/28	150	42	54.8 (4.1)	82
Total	1819	501	18.2 (0.9)	330

Table 4. Chinook salmon mid eye to fork length (mm) by age and sex, 2003, Tozitna River, Alaska.

Age	Sex	N	Mean	SE	Range
1.1	Male	2	378	48	320-425
	Female	0	-	-	-
1.2	Male	139	518	4.3	400-755
	Female	0	-	-	-
1.3	Male	231	703	3.5	500-850
	Female	28	778	7.1	725-895
1.4	Male	37	793	15	440-975
	Female	62	859	5.2	780-940
1.5	Male	1	805	-	805
	Female	1	865	-	865

Table 5. Summer chum salmon escapement age composition by stratum and sex, Tozitna River, Alaska, 2003. Standard error in parentheses.

		Brood Year and Age										
Strata	Run	Sex	2000		1999		1998		1997		Total	
			0.2		0.3		0.4		0.5		N	%
			N	%	N	%	N	%	N	%	N	%
6/26-7/7	14	Males	0	0.0	6 (16.3)	60.0	4(16.3)	40.0	0	0.0	10	100
		Females	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
		Subtotal	0	0.0	6 (16.3)	60.0	4 (16.3)	40.0	0	0.0	10	100
7/8-7/14	722	Males	1 (1)	0.6	83 (3.9)	51.2	15 (3.5)	9.3	4 (1.9)	2.5	103	64
		Females	0	0.0	49 (4.9)	30.2	7 (4.2)	4.3	3 (2.9)	1.9	59	36
		Subtotal	1 (.6)	0.6	132 (3.1)	81.4	22 (2.7)	13.6	7 (1.6)	4.4	162	100
7/15-7/21	2636	Males	1 (.7)	0.4	119 (3.3)	50.9	22 (2.9)	9.4	6 (1.6)	2.6	148	63
		Females	1 (1.2)	0.4	71 (4.1)	30.3	13 (3.9)	5.6	1 (1.2)	0.4	86	37
		Subtotal	2 (.6)	0.8	190 (2.6)	81.2	35 (2.3)	15.0	7 (1.1)	3.0	234	100
7/22-7/28	5115	Males	1 (1)	0.7	88 (3.3)	59.0	11 (3.1)	7.4	1 (1)	0.7	101	68
		Females	1 (2.1)	0.7	45 (3.5)	30.2	2 (2.9)	1.3	0	0.0	48	32
		Subtotal	2 (1)	1.4	133 (2.5)	89.2	13 (2.3)	8.7	1(.7)	0.7	149	100
Subtotal		Males	3 (1.1)	0.5	296 (3.4)	53.3	52 (3.2)	9.4	11 (1.3)	2.0	362	66
		Females	2 (1.7)	0.4	165 (3.9)	29.7	22 (3.4)	4.0	4 (1.1)	0.7	193	34
Total	8487		5 (1)	0.9	461 (2.6)	83.0	74 (2.4)	13.4	15 (1.0)	2.7	555	100

Table 6. Proportion and estimated number of female summer chum salmon, Tozitna River, Alaska, 2003. Standard error in parentheses.

Strata	Run	N	Percent Female	Estimated # Females
6/26-7/7	14	10	0.0	0
7/8-7/14	722	162	36.4 (1.8)	263
7/15-7/21	2636	234	36.8 (0.9)	969
7/22-7/28	5115	149	32.2 (0.7)	1648
Total	8487	555	33.9 ^a (0.5)	2880 ^a

^aCalculations were determined with partial escapement data because of high stream discharge.

Table 7. Summer chum salmon mid eye to fork length (mm) by age and sex, 2003, Tozitna River, Alaska. SE = Standard Error

Age	Sex	N	Mean	SE	Range
0.2 ^a	Male	3	560	8.7	545-575
	Female	2	530	15	515-545
0.3 ^a	Male	296	575	1.6	510-655
	Female	165	550	2.1	500-620
0.4 ^a	Male	52	608	4.4	525-675
	Female	22	590	5.4	540-640
0.5 ^a	Male	11	626	9.6	575-675
	Female	4	605	9.6	580-620

^aCalculations were determined with partial escapement data because of high stream discharge.

Table 8. Comparison of preliminary Chinook salmon escapement age composition by sex at the East Fork (EF) of Andreafsky River, Gisasa River, Henshaw Creek, and the Tozitna River, Alaska, 2003.

Location	Yukon River (km)	N	Sex	Brood year and Age					Total
				2000	1999	1998	1997	1996	
				%	%	%	%	%	%
EF Andreafsky Weir	167 ^a	533 ^d	Males	0.4	12.7	33.2	6.0	0.0	52.3
			Females	0.0	3.2	17.3	26.1	1.1	47.7
			Subtotal	0.4	15.9	50.5	32.1	1.1	100.0
Gisasa Weir	818 ^b	472 ^d	Males	0.2	5.5	51.3	4.9	0.0	61.9
			Females	0.0	0.0	18.2	18.8	1.1	38.1
			Subtotal	0.2	5.5	69.5	23.7	1.1	100.0
Henshaw Weir	818 ^b	304 ^d	Males	1.6	19.4	35.5	4.3	0.0	60.9
			Females	0.0	0.0	8.6	28.9	1.6	39.1
			Subtotal	1.6	19.4	44.1	33.2	1.6	100.0
Tozitna Weir	1096 ^c	501 ^e	Males	0.4	27.7	46.1	7.4	0.2	81.8
			Females	0.0	0.0	5.6	12.4	0.2	18.2
			Subtotal	0.4	27.7	51.7	19.8	0.4	100.0

^aKilometers from mouth of the Andreafsky River to mouth of the Yukon River.

^bKilometers from the mouth of the Koyukuk River to the mouth of the Yukon River.

^cKilometers from the mouth of the Tozitna River to the mouth of the Yukon River.

^dPreliminary escapement age data from ADF&G, 2003.

^ePreliminary escapement age data from BLM, 2003.




Figure 1. Location of the Tozitna River weir, Alaska, 2003.




Figure 2. Chinook salmon daily counts with quartiles shown (25, 50, and 75%) of cumulative escapement for the period 26 June – 26 July, 2003, Tozitna River, Alaska.




Figure 3. Daily discharge (m³/s) for the period 21 June – 11 August 2003, Tozitna River, Alaska.




Figure 4. Map showing the locations of four weirs in Alaska monitoring Chinook salmon escapement on Yukon River tributaries: East Fork of the Andreafsky River, Henshaw Creek, Gisasa River, and the Tozitna River. Map is adapted from Holder and Senecal-Albrecht (1998).

The U.S. Fish and Wildlife Service, Office of Subsistence Management conducts all programs and activities free from discrimination on the basis of sex, color, race, religion, national origin, age, marital status, pregnancy, parenthood, or disability. For information on alternative formats available for this publication please contact the Office of Subsistence Management to make necessary arrangements. Any person who believes she or he has been discriminated against should write to: Office of Subsistence Management, 3601 C Street, Suite 1030, Anchorage, AK 99503; or O.E.O., U.S. Department of Interior, Washington, D.C. 20240.